

TITLE OF THE INVENTION

New Metalloproteases of the Neprilysin Family

BACKGROUND OF THE INVENTION

5 Peptides are used by cells from yeast to mammals to elicit physiological responses. The use of peptides as messengers usually involves the following steps: 1) production and release of the peptide by a specific cell, 2) interaction of the peptide with a receptor on the surface of the target cell, and 3) degradation of the peptide to terminate its action. The first and last steps of this scheme require the participation of

10 proteases/peptidases. There is increasing evidence that membrane-associated zinc-metallopeptidases play important roles in both of these steps. Although activation of prohormone precursors into bioactive peptides is generally performed by proteases of the subtilisin family located in the *Trans*-Golgi Network or in secretory granules of the cell (for a review see: (Seidah and Chrétien, 1995)) a few peptides need a final

15 processing step. This step involves the action of membrane-associated zinc-metallopeptidases. Two cases are particularly well documented: angiotensin-converting enzyme (ACE) which cleaves inactive angiotensin I into angiotensin II (Corvol and Williams, 1997) and endothelin-converting enzymes (ECEs) which cleave isoforms of big endothelins into endothelins (Turner, 1997a). In addition to their role

20 in peptide activation, cell surface zinc-metallopeptidases have also been implicated in the termination of the peptidergic signal by breaking down the active peptides into inactive fragments. One of the best known of these peptidases is probably Neutral Endopeptidase-24.11 (Neprilysin, NEP) that has been implicated in the physiological degradation of several bioactive peptides (Kenny, 1993). Interestingly, NEP and the

25 ECEs show significant structural similarities and appear to be members of a family of peptidases that also includes PEX, a newly discovered and not yet characterized peptidase, and the KELL blood group protein (Turner and Tanzawa, 1997b). Because of their important role as regulators of bioactive peptide activity, these enzymes (more specifically NEP and the ECEs) have been identified as putative targets for therapeutic

30 intervention, similar to the way ACE inhibitors are used to control blood pressure. The recent discovery of PEX, another member of the family, which appears to be involved in phosphate homeostasis, raised the possibility that other yet unknown members might exist.

35 Members of the NEP-like family are type II membrane proteins consisting of three distinct domains: a short NH₂-terminal cytosolic sequence, a single transmembrane region, and a large extracellular or ectodomain responsible for the catalytic activity of the enzyme. There are potential N-glycosylation sites and cysteine residues that are involved in disulfide bridges stabilizing the conformation of the active enzyme. These enzymes are metalloenzymes with a Zn atom in their active site. As

such, they belong to the zincin family of peptidases which is characterized by the active site consensus sequence HEXXH (Hooper, 1994), where the two histidine residues are zinc ligands. In members of the NEP-like family of peptidases, the third zinc ligand is a glutamic acid residue located on the carboxy-terminus side of the consensus sequence. This characteristic puts them in the gluzincin sub-family (Hooper, 1994). The model enzyme for gluzincins is thermolysin (TLN) a bacterial protease whose 3D structure has been determined by X-ray crystallography (Holmes and Matthews, 1982). The active site of NEP has been extensively studied by site-directed mutagenesis and several residues involved in zinc binding (Devault et al., 1988b; Le Moual et al., 1991; Le Moual et al., 1994), catalysis (Devault et al., 1988a; Dion et al., 1993), or substrate binding (Vijayaraghavan et al., 1990; Beaumont et al., 1991; Dion et al., 1995; Marie-Claire et al., 1997) have been identified (for a recent review see Crine et al., 1997).

SUMMARY OF THE INVENTION

Here, we developed an RT-PCR strategy to look for other members of this important family of peptidases. This strategy allowed the molecular cloning and characterization of three additional NEP-like (NL) metallopeptidases (called NL-1, NL-2 and NL-3). Knowledge obtained through these studies allows the generation of reagents (nucleic acid probes and primers, antibodies and active recombinant enzymes) for further biochemical characterization of these enzymes and their pattern of expression and will greatly help the rational design of specific inhibitors that could be used as therapeutic agents.

Accordingly, the present invention relates to the following products:

- A. Degenerate primers for screening new NEP-related enzymes;
- B. NL-1, NL-2 and NL-3 proteins as NEP-related enzymes;
- C. Nucleic acids encoding these enzymes;
- D. Antibodies directed against the enzymes;
- E. Recombinant vectors comprising the nucleic acids encoding the enzymes and hosts transformed therewith;
- F. Fragments of the nucleic acids useful as probes or primers to hybridize and detect the presence of an NL-1, NL-2 and NL-3 genes, or to hybridize and amplify and produce gene fragments;
- G. Soluble forms of NL-1, NL-2 and NL-3; and
- H. Nucleic acids comprising the N-terminal part of NL-1 or NL-2 which terminates with a sequence encoding a furin recognition site, such nucleic acids being useful for making a fusion protein with the ectodomain of any protein of interest, and for releasing a soluble form of that protein of interest (containing the ectodomain) in the medium.

Also the present invention relates to the following methods:

- A. A method for screening NEP-related enzymes that make use of degenerate primers or probes selected from a region of NEP family members in a highly conserved region, namely around the zinc-binding sites; and
- 5 B. A method for producing NL-1, NL-2 or NL-3 that includes the steps of culturing the above recombinant host and recovering NL-1, NL-2 and NL-3 gene products therefrom.

The present invention will be described hereinbelow by referring to specific embodiments and appended figures, which purpose is to illustrate the invention rather than to limit its scope.

In the first section, general procedures leading to the identification and localization of NL-1, NL-2 and NL-3 are given. In the second section, slightly different procedures are given for completing or reiterating the work performed on NL-1.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Amino acid sequence comparison of human NEP, PEX, KELL and ECE1 peptidases. Amino acid sequences in boxes are those used to design the oligonucleotide primers. Numbers and arrows under the sequences identify the primer and its orientation.

Figure 2: Sequences of the oligonucleotide primers used in the PCR reactions.

Figure 3: Nucleotide and amino acid sequence of the mouse NL-1 cDNA. The sequence of the DNA fragment obtained by PCR is in brackets.

Figure 4: Partial nucleotide and amino acid sequence of the human NL-2 cDNA. The sequence of the DNA fragment obtained by PCR is in brackets.

Figure 5: Partial nucleotide and amino acid sequence of the human NL-3 cDNA.

Figure 6: Amino acid sequence comparison of NEP, NL-1, NL-2 and NL-3 peptidases.

Figure 7: *In situ* hybridization of mouse testis sections using NL-1 as a probe.

Figure 8: *In situ* hybridization of mouse sections using mouse NL-3 as a probe.

Figure 9: *In situ* hybridization of mouse spinal chord sections

Figure 10: Expression of NL-1 in mammalian cells.

Figure 11: Activity of recombinant soluble NL-1.

Figure 12: Expression of a soluble form of NL-3 using NL-1 amino-terminal domain.

DESCRIPTION OF THE PREFERRED EMBODIMENTS OF THE INVENTION

SECTION 1)

5 MATERIALS AND METHODS

DNA and RNA manipulations

All DNA manipulations and Northern blot analysis were performed according to standard protocols (Ausubel et al., 1988; Sambrook et al., 1989).

mRNA purification and cDNA synthesis

10 mRNAs were prepared from mouse testis using Quick Prep Micro mRNA purification kit (Pharmacia Biotech). Purified mRNAs were kept at -70° until ready used. First strand cDNA was synthesized from 1µg of mRNA using the First-Strand cDNA synthesis kit (Pharmacia Biotech). The human testis cDNA library was obtained from Clontech.

15 Polymerase chain reaction protocol

PCR was performed in a DNA thermal cycler with 5 µl of cDNA template and 1 µl of Taq DNA polymerase in a final volume of 100 µl, containing 1 mM MgCl₂, 2 µM of each primer oligonucleotide, 20 µM of each dNTP and 5% DMSO. Cycling profiles included an initial denaturation step of 5 min at 94°C, followed by 30 cycles of 1 min at 94°C, 1 min at 40°C and 1.5 min at 72°C. A final extension step was performed at 72°C for 10 min. The amplified DNA was loaded on a 2% agarose gel and visualized by staining with ethidium bromide. Fragments ranging in size between 500-700 bp were cut and eluted from the gel. If needed, a second round of PCR was done with nested oligonucleotide primers, using 10 µl of the first PCR reaction, or of the eluted band cut from the agarose gel. Resulting fragments were ligated in pCR2.1 vector (Invitrogen) according to the distributor's recommendations. DH5α *Escherichia coli* cells were transformed with the ligation mixture and grown on 2YT plates in the presence of kanamycin. Plasmids were prepared from resistant cells and sequenced.

25 In situ hybridization on mouse tissues and chromosomal localization of human genes

30 *In situ* hybridization on whole mouse slices or isolated tissues was performed as described previously (Ruchon et al., 1998).

To determine the chromosomal localization of human NL-2 and NL-3 genes, a technique for mapping genes directly to banded human chromosomes was used. Metaphase chromosomes were obtained from lymphocytes cultured from normal human peripheral blood. Cells were synchronized with thymidine and treated with 5-bromodeoxyuridine (BrdU) during the last part of the S phase to produce R-banding. Biotin-labeling of the probe was done by nick-translation (Bionick, BRL) and the probe was visualized by indirect immunofluorescence.

35 Antibody production

prepare extracts of cellular proteins or culture media were also described in previous papers (Devault *et al.*, 1988a; Lemay *et al.*, 1989). The presence of NLs in these extracts was monitored by immunoblotting using specific antibodies.

5 Extracts of cellular proteins and culture media were assayed for enzymatic activity. Two tests were performed. The first used [³H]-Tyr-(D)Ala₂-Leu-enkephalin as substrate and was performed according to Lemay *et al.*, (1989). The second used bradykinin as substrate and was performed as described by Raut *et al.* (1999).

RESULTS

10 Cloning of NL-1, a new member of the NEP family

The molecular cloning in the past few years of ECEs, PEX and KELL showed that all these proteins have between 50 and 60% similarity with NEP. This observation led us to believe that these peptidases are part of an extended family and that there could be still additional members to be discovered. To test this hypothesis, we aligned the amino acid sequences of the members of the NEP-like family and designed degenerate oligonucleotide primers to be used in RT-PCR reactions (Figure 1 and 2). These primers were located on either side of the HEXXH consensus sequence for zincins. Because they are highly degenerate, primers 1 and 2 were each subdivided into two pools, 1A-1B, and 2A-2B, respectively (Figure 2). Any PCR amplified DNA fragment that corresponds to a peptidase of the family should normally contain the consensus sequence and be easily recognized by sequencing of the cloned fragments. Using this strategy, we first performed PCR reactions with primer pairs 1A-3 and 1B-3. The amplified DNA migrates mostly as a smear starting at around 700 bp and going down to 100 bp. As the expected fragments should be around 550 bp, we isolated from the gel the section corresponding to DNA fragments longer than 500 bp. A second round of PCR reactions was performed with both crude PCR products of the first reaction and isolated DNA bands, using primers 2A-3 and 2B-3. The expected 296 bp fragment was seen on the gel (not shown).

Cloning of these DNA fragments generated approximately 350 clones, of which 30 44 were sequenced. Nine of these had no inserts or corresponded to sequences not related to the NEP family, 24 corresponded to NEP, 3 to PEX, and 8 corresponded to one putative new member of the family, since they all contained the HEXXH consensus sequence for zincins and showed 65% homology with mouse NEP (in boxes Figure 3). This fragment was then used to screen a mouse testis cDNA library, and allowed us to isolate a complete cDNA of 2592 nucleotides (Figure 3). The identity of this sequence with other members of the family is presented in Table I. This new member was called NL-1, for NEP-like peptidase 1.

Cloning of NL-2 and NL-3.

A strategy similar to that described for amplification of enzymes of the NEP family from mouse testis cDNAs was used with a human testis cDNA library using two different oligonucleotide primers. This time, DNA fragments of approximately 900 bp were obtained and cloned. Ten clones were sequenced, revealing the presence of NEP and two new peptidases of the family that we have called NL-2 and NL-3.

The NL-2 PCR fragment was 879 nucleotides in length and encoded a 293 amino acid residue segment probably located in the carboxy-terminal domain of this putative peptidase (in brackets Figure 4). This PCR fragment was then used to screen a lambda gt10 human brain cDNA library. It allowed the isolation of other cDNA fragments which overlap partially with the NL-2 PCR fragment. Fusion of these lambda clones and the PCR fragment resulted in an open reading frame of 770 amino acid residues. The use of 5' RACE protocols with human testis cDNA libraries allowed completion of the sequence of NL-2 ORF (Figure 4). This ORF codes for a putative protein that is about 80% identical to the mouse NL-1 protein (Figure 6). Across species, members of the NEP, PEX, ECEs sub-families have highly conserved sequences (more than 94% identity). Although a sequence identity of about 80% only exists between the novel human protein and mouse NL-1, these proteins share unique characteristics that make possible the fact that NL-2 protein may be the human homologue of NL-1. The identity of NL-2 with other members of the family is presented in Table I.

The 879 bp PCR fragment encoding NL-3 showed an open reading frame of 293 amino acid residues (Figure 5, in brackets). Sequence analysis of NL-3 showed that it was 94.2 % identical to an EST sequence from mouse embryonic tissue present in publicly accessible DNA data banks. This mouse EST sequence, commercially available from American Tissue and Cell Culture (ATCC), had been obtained previously by our laboratories.

Since Northern blot analysis of human tissues with the NL-3 PCR fragment showed the expression of this protein in spinal chord (see below), the same PCR DNA fragment was used to screen by hybridization a human spinal chord cDNA library constructed in phage λ vectors. One clone contained a full-length ORF of 752 amino acid residues that encompassed the 293 amino acid residue ORF of the PCR fragment. Further probing, cloning and sequencing lead to the obtention of NL-3 full sequence, shown in Figure 5.

Figure 6 presents a comparison of the amino acid sequence of the new NEP-like enzymes and Table I shows the extent of identity between members of the family.

Cellular distribution of NL-1, NL-2 and NL-3 peptidases

Determining the tissue distribution of NL-1, NL-2 and NL-3 may provide clues to identify the peptidergic systems in which they are involved. It will be particularly

interesting to compare the tissue distribution of these peptidases with that of NEP and the ECEs to determine whether or not the physiological functions of NL-1 and/or NL-2 and/or NL-3 may overlap those of NEP and/or ECEs.

5 *In situ* hybridization (ISH), using our mouse cDNA, was used to determine the spatial and temporal expression of NL-1 during mouse development, as done previously for PEX (Ruchon et al., 1998)). Serial sections of whole foetal (12, 15 and 19 dpc) and adult mice (1, 3 and 6 days old) were hybridized with an [³⁵S]-labeled RNA probe. Figure 7 shows a section of mouse testis which was the only tissue identified to express NL-1 by this technique. Cells of seminiferous tubules are specifically labeled but spermatids located near the center of the tubule showed strongest labeling. These cells are in the last stage of maturation into spermatozooids. The presence of NL-1 in testis has now been confirmed by Northern analysis of mouse tissues (see Fig. 10). Other tissues express NL-1, when analyzed by RT-PCT, which is a more sensitive assay (not shown).

15 A similar approach was used to determine the localization of NL-3 using the mouse EST obtained from ATCC. Figure 8 shows sections of whole mouse at 17 days of embryonic development and 4 days post-natal. Several tissues are expressing this putative peptidase including brain, where it is associated with neurons (Figure 9), spinal chord, liver, spleen and bones. Labeling was stronger in bones from *Hyp* mouse, an animal model for hypophosphatemic rickets (Figure 8). In bones, NL-3 was found to be expressed by osteoblasts (not shown).

20 Northern blotting experiments were performed on several tissues with NL-2 and NL-3 probes. A Human Multiple Tissues Northern Blot (Clontech) was hybridized with specific probes. A single RNA band of approximately 4.0 kb was revealed by the probe for NL-2. Expression of NL-2 is restricted to brain and spinal cord (not shown). However, RT-PCR has shown the presence of this enzyme in testis (not shown).

25 A single RNA band of approximately 3.0 kb was detected with the specific probe for NL-3 (not shown). NL-3 expression was observed mainly in ovary, spinal cord and adrenal gland.

30 Chromosomal localisation of the human gene for NL-2 and NL-3

As a mean to get clues on the function of the new metallopeptidases in vertebrates, we have localized the new cDNAs on human chromosomes, in order to look for a possible link between the gene locus and mapped genetic diseases in humans. To do so, we have mapped the NL-2 and NL-3 genes by high-resolution fluorescence *in situ* hybridization (FISH). NL-2 was localized to chromosome band 1p36. Consistent with the cellular distribution of NL-2 in humans, genetic diseases of the CNS such as dyslexia, neural tube defect, neuroblastoma, neuronal type of Charcot-Marie-Tooth disease have all been mapped in this region and represent potential targets for a role of NL-2 in humans. NL-3 was localized to chromosome band

35

2q37. Consistent with a role of NL-3 in bones, a form of Albright hereditary osteodystrophy was mapped to the same chromosomal locus (Phelan et al., 1995).

In view of the foregoing, NL-2 and NL-3 are metallopeptidases that are assumed to be immediately useful as markers for a disease or disorder associated with human chromosomal locus 1p36 and 2q37, respectively. Their localization on a chromosome band associated with known diseases suggests that they may be expressed or co-expressed with one or more genes, as a cause or a consequence of disease development. It is possible that these enzymes are up or down regulated, alone or along with other genes involved in a disease. Therefore, antibodies or other ligands specific to NL-2 or NL-3 may be used for a diagnostic purpose, as well as primers or probes in diagnostic assays using nucleic acid hybridization or amplification techniques. Otherwise, primers or probes directed against the nucleic acids of NL-2 and NL-3 would be useful to map the mutations of a gene located in close proximity and involved in the disease. Therefore, no matter which exact function NL-2 and NL-3 gene products have, their chromosomal localization provides one diagnostic utility. This localization as well as the tissular distribution provide information as to the disease and tissue to be investigated to elucidate the exact function of these enzymes.

NL-1 resembles NL-2, sharing with the latter about 80% homology in the amino sequence and sharing structural characteristics such as the furin recognition sequence located at the proximal end of the ectodomain. NL-2 might be the human homologue of mouse NL-1. If such was the case, these two proteins would have a substantial degree of divergence and, maybe, different profiles of activity varying from one species to another.

Chromosomal localization of NL-1 was determined in mouse genome by Single Strand Conformational Polymorphism (SSCP) in collaboration with The Jackson Laboratory Backcross DNA Panel Mapping Resource. NL-1 was localized to the distal region of mouse chromosome 4 which corresponds to human chromosome region 1p36 where is located NL-2 gene. This reinforces our hypothesis that NL-1 and NL-2 are species variants.

Production of antibodies against NLs

Antisera collected from injected animals were first tested by immunoblotting on GST-antigen fusion proteins produced in *E. coli*. Antiserum from one rabbit recognized the NL-1-related polypeptide and antisera from one mouse and one rabbit reacted with the NL-3-related polypeptide (results not shown). The anti NL-1 antiserum and the mouse anti NL-3 antiserum, which appeared more specific than the rabbit antiserum, were next tested by immunoblotting on extracts of proteins and culture media from cells expressing NL-1 or NL-3 (see below).

Expression of NL-1 in CHO cells

The cDNA encoding the full-length NL-1 protein was cloned in the mammalian

expression vector pcDNA3-RSV and transfected in CHO cells. Stable cell lines were established by selection with the drug G418 and tested by immunoblotting for the presence of NL-1.

5 Small amounts of NL-1 were found in the extracts of transfected CHO cells (results not shown). This intracellular species was sensitive to endo H digestion, indicating that the sugar moiety was not mature and suggesting ER localization (results not shown). The culture medium of transfected CHO cells showed the presence of soluble NL-1 (Figure 10). This extracellular species was resistant to endo H suggesting true transport through the late secretory pathway. The cDNA sequence of NL-1
10 predicts a type II transmembrane protein. The mechanism by which NL-1 is transformed into a soluble protein is not known presently. However, examination of the amino acid sequence revealed the presence of a putative furin cleavage site from residue 58 to 65 (Figure 3). A similar site is present in NL-2 sequence.

The soluble form of NL-1 was assayed for activity using [³H]-Tyr-(D)Ala₂-Leu-
15 enkephalin and bradykinin as substrates. Figure 11 shows that NL-1 can degrade the enkephalin substrate ($K_m = 18 \pm 10 \mu\text{M}$) and that this activity can be inhibited by phosphoramidon ($IC_{50} = 0.9 \pm 0.3 \text{ nM}$) and thiorphan ($K_m = 47 \pm 12 \text{ nM}$), a general inhibitor of enzymes of the NEP family. Bradykinin is also a substrate for NL-1 (not shown).

Use of NL-1 amino-terminal domain to promote secretion

20 The observation that NL-1 ectodomain was secreted, possibly through cleavage of the transmembrane segment by furin, raised the possibility to promote secretion of exogenous proteins that could be spliced to NL-1 amino-terminal domain (from initiator methionine to the furin site). To test this hypothesis, the ectodomain of NL-3 (from the third cysteine to the end) was spliced to NL-1 amino-terminal domain using a PCR
25 strategy and the recombinant DNA cloned in expression vector pRcCMV. The fusion protein was expressed by transfection of the vector in COS-1 and HEK 293 cells. The culture media of transfected cells was analyzed by immunoblotting using the mouse antiserum against NL-3. Figure 12 shows the presence of NL-3 in the spent culture media of both COS-1 and HEK 293 cells. This result shows that NL-1 amino-terminal
30 domain can be used to promote secretion of exogenous proteins.

The soluble form of NL-3 was assayed for activity using [³H]-Tyr-(D)Ala₂-Leu-enkephalin as substrate. No activity was found.

The previous experiment showed that it was possible to use the amino-terminal domain of NL-1 to promote secretion of an otherwise membrane attached protein
35 ectodomain. To verify whether the same strategy could be used to promote secretion of small peptides, a PCR strategy was used to splice human β -endorphin to the amino-terminal domain of NL-1 and the recombinant DNA was cloned in vector pRcCMV. The fusion protein was expressed by transfection of the vector in COS-1 and HEK 293 cells. The culture media of transfected cells was collected 48h after transfection and

the peptides purified as described previously (Noël et al., 1989). The presence of β -endorphin in the extracts was detected by radioimmunoassay. The results showed that both COS-1 and HEK 293 cells produced approximately 100 pg of β -endorphin per ml of culture medium. Therefore, the N-terminus of LN-1 and NL-2 which ends with a furin-recognition site will be useful to produce the soluble form of a protein of interest.

SECTION 2)

MATERIALS AND METHODS

10 DNA manipulations

All DNA manipulations, phage library screening, and plasmid preparations were performed according to standard protocols (Ausubel 1988; Sambrook 1989). Site-directed mutagenesis was performed using a PCR-based strategy as described previously (Le Moual 1994).

15

mRNA purification and RT-PCR protocol for identification of new members of the neprilysin family

mRNAs were prepared from mouse testis using Quick Prep Micro mRNA purification kit (Pharmacia Biotech). First strand cDNA was synthesized from 1 μ g of mRNA using the First-Strand cDNA synthesis kit (Pharmacia Biotech).

20

Two sense primers, oligonucleotide 3817 (5'-TGGATGGAT/CGA/CIGG/AIACIA/CA-3') and oligonucleotide 3719 (5'-A/GTIGTITTT/CCCIGCIGGIA/GT/AIC/TTG/CCA-3') corresponding respectively to amino acid residues 459 to 465 and 552 to 560 of NEP sequence, and one antisense primer, oligonucleotide 3720 (5'-AIICCICCIA/TC/TA/GTCIGCIG/AC/TA/GTTT/CTC-3') corresponding to amino acid residues 646 to 654 (see Fig. 1 and 2), were synthesized. PCR was performed with 5 μ l of cDNA template and 1 μ l of Taq DNA polymerase in a final volume of 100 μ l, containing 1 mM MgCl₂, 2 μ M of each oligonucleotide 3817 and 3720, 200 μ M of each dNTP and 5% DMSO. Cycling profiles included an initial denaturation step of 5 min at 94°C, 30 cycles of 1 min at 94°C, 1 min at 40°C and 1.5 min at 72°C, and a final extension step at 72°C for 10 min. One half of the amplified DNA was fractionated on a 2% agarose gel and fragments ranging in size between 500-700 bp were purified and resuspended in a final volume of 50 μ l. A second round of PCR was done with primers 3719 and 3720, using as template either 10 μ l of the first PCR reaction or 5 μ l of the purified fragments, and the new PCR products were ligated in pCR2.1 vector (Invitrogen). Several identical clones corresponded to a potential new member of the NEP family. We called this member NL1 for NEP-like 1.

25

30

35

Cloning of full-length NL1 cDNA

The cloned NL1 PCR fragment was used as probe to screen a mouse testis λ Uni-ZAP™XR cDNA library (Stratagene). Twelve out of a hundred positive phages were plaque purified and subcloned into pBS SK vector (Stratagene). As the longest clone analyzed presented an incomplete ORF (pBS-NL1A), 5'RACE with primers located in

5 vector (5'-TAGTGGATCCCCCGGGCTGCAG-3', sense primer) and NL1 (5'-ACCAAACCTTTCCTGTAGCTCC-3', antisense primer, nt 1303 to 1324 of NL1); was subsequently performed on the DNA of the remaining semi-purified positive clones. Amplification was performed with 1 μ l of Vent polymerase in a final volume of 100 μ l containing 50 ng of DNA, 4 mM of MgSO₄, 1 μ M of each oligonucleotide, 200 μ M of each dNTP and 10% DMSO. Cycling parameters included an initial denaturation step

10 of 1 min at 94°C, 25 cycles of 30 sec at 94°C, 30 sec at 60°C and 1 min at 72°C, and an incubation of 10 min at 72°C. A PCR fragment of the expected length was subcloned into pCR2.1 vector (clone pCR-NL1A), but sequencing revealed no initiator ATG codon. A nested 5'RACE was then performed on mouse testis cDNA using the

15 Marathon Ready cDNA kit (Clontech) with sense oligonucleotides AP1 and AP2 (from the kit) and NL1 antisense oligonucleotides 5'-CCTGAGGGCTCGTTTTACAACCGTCCT-3' (nt 503 to 529 of NL1) and 5'-CTCATCCCAGGAGAAGTGTAGCAGGCT-3' (nt 475 to 502 of NL1) as recommended by the supplier. The resulting fragment was cloned into pCR2.1 vector (pCR-NL1B).

20 Since only ten bp were missing for the initiator ATG codon, we reconstructed the 5' end of the cDNA by PCR-amplifying clone pCR-NL1A with sense primer 5'-CCACCATGGTGGAGAGAGCAGGCTGGTGTCCGGAAGAAG-3' (nt 332 to 364 of NL1; the 10 missing nucleotides are underlined) and antisense primer 5'-ACCAAACCTTTCCTGTAGCTCC-3' (nt 1303 to 1324 of NL1) using Vent polymerase

25 as described above. The DNA fragment was then inserted into pCR2.1 (clone pCR-NL1C). The entire ORF was reconstituted following digestion of pBS-NL1A and pCR-NL1C with *Eco*RI and *Pf*MI. The 5' end of NL1 cDNA was excised from pCR-NL1C and ligated into pBS-NL1A at the corresponding sites, resulting in plasmid pBS-NL1B.

For expression studies, a *Bam*HI/*Ap*al fragment generated out of pBS-NL1B, corresponding to the full length cDNA of NL1, was inserted into pCDNA3/RSV [18] vector.

30

Production of polyclonal antibodies

A plasmid for the production in *Escherichia coli* of a GST fusion protein with NL1 was constructed using pGEX-4T-3 expression vector (Pharmacia Biotechnologies). A 255 bp fragment from NL1 was amplified by PCR with Vent polymerase using sense primer 5'-GCTACGGGATCCCGTGGCCACTATGCTTAGGAA-3' (nt 1139 to 1158) and antisense primer 5'-CGATTGCTCGAGTGGGAACAGCTCGACTTCCA-3' (nt 1377 to 1396). Both pGEX-4T-3 and the PCR product were digested with *Bam*HI and *Xho*I

35

and ligated. The recombinant protein was produced and purified according to the supplier's instructions. Five weeks old female balb/c mice were immunized at monthly intervals for 3 months with 20 µg of the recombinant NL1 fragment in Freund's adjuvant and antisera were subsequently collected.

5

Cell culture and transfection

Human Embryonic Kidney (HEK 293) cells were maintained in DMEM medium containing 10% fetal bovine serum (FBS), and supplemented with penicillin at 60 µg/ml, streptomycin at 100 µg/ml and fongizone at 0,25 µg/ml. Transfections of cells with appropriate plasmids were performed by the calcium/phosphate-DNA co-precipitation method (Chang 1987). To establish permanent cell lines, G418 selection was initiated 48 h after the transfections at 400 µg/ml for 12 days and gradually decreased at 100 µg/ml.

LLC-PK₁ cells transfected with pRcCMV-sNEP were maintained as described previously (Lanctôt 1995).

15

Immunoblot analysis

For immunoblot analysis, cells were incubated for 16 h in synthetic DMEM medium containing 2mM sodium butyrate. Cellular proteins were solubilized as previously described (Dion 1995). Secreted proteins recovered in culture media were concentrated approximately 10 fold by ultrafiltration. Immunoblot analysis were performed using the NEN Renaissance kit with the polyclonal antibody specific to NL1 or the α1-antitrypsin inhibitor antibody (Calbiochem; LaJolla, CA) followed by the appropriate horseradish peroxidase-conjugated IgG (Vector Laboratories).

20

For the glycosylation studies, proteins were incubated with endoglycosidase H (endoH) or peptide:N-glycosidase (PNGaseF) as suggested by the distributor (NEB).

25

Enzymatic activity assays

NL1 activity was monitored and compared to sNEP activity using (Tyrosyl-[3,5-³H])(D-Ala₂)-Leu₅-enkephalin (50 Ci/mmol) (Research Products International Inc.), as already described (Dion 1995; Devault 1988). *K_m* values were determined by the isotope-dilution method. The inhibitory effects of phosphoramidon and thiorphan were also assessed as previously described (Dion 1995).

30

HPLC analysis of the hydrolysis of Leu-enkephalin

Five µg of Leu₅-enkephalin were incubated at 37°C for one hour in 50mM MES, pH 6.5, with concentrated culture medium of HEK 293 cells expressing NL1 (~300 µg of total proteins) or LLC-PK₁ cells expressing sNEP (~30 µg of total proteins), in absence or presence of 0.1 mM phosphoramidon. Hydrolysis products were separated by

35

FOR "SECRET" 6666666666

reversed-phase HPLC as described previously [23]. Tyr-Gly-Gly and Phe-Leu were both identified by elution profiles of synthetic marker peptides.

Northern blot analysis

5 A mouse multiple tissue poly(A)⁺ mRNA blot (Clontech) was hybridized with a [³²P]dCTP random primer labelled probe in ExpressHyb solution (Clontech). The blot was washed according to the manufacturer's recommendations and exposed to Fuji RX film for 7 days at -80°C with intensifying screens.

10 RT-PCR screening of mouse tissues

First strand cDNA synthesis was performed with 1 µg of total RNA from mouse tissues and oligo(dT) as primer, using Gene Amp RNA PCR Core Kit (Perkin Elmer). For the PCR reactions, primers 5'-TGGCGAGAGTGTGTCAGCTATGTC-3' and 5'-CTTCCAAAATGTAGTCAGGGTAGCCAATC-3' were used with Taq polymerase. One 15 tenth of the PCR products were visualized on a 4% agarose gel.

In situ hybridization

To construct a plasmid for the synthesis of cRNA probes for ISH, pCR-NL1A was used as template to amplify a 452 bp fragment by PCR with sense primer 5'- 20 GGAGCCATAGTGA CTCTGGGTGTC-3' (nt 416 to 439) and antisense primer 5'-GACGCTCAGCAGGGGCTCAGAGTC-3' (nt 842 to 865). The amplification product was inserted into pCRII vector (Invitrogen). Synthesis of riboprobes and protocols for ISH were as described previously (Ruchon 1998).

RESULTS

25

Cloning and sequence analysis of mouse NL1 cDNA

In order to isolate cDNAs for new members of the NEP family, we developed an RT-PCR strategy based on fact that NEP, ECE-1 and PHEX share regions of significant sequence identity. Following RT-PCR on testis mRNAs with nested primers, 30 a DNA fragment of approximately 300 bp was amplified. This DNA fragment was cloned and the plasmids from 24 independent colonies were sequenced: 3 clones had no insert, 4 clones had DNA fragments not related to the NEP family, 7 clones had sequences corresponding to mouse NEP and 3 clones had sequences corresponding to mouse PHEX, showing that our approach efficiently allowed the identification of 35 members of the family. Moreover, 7 identical clones had a new cDNA presenting sequence similarities to members of the NEP family. The full-length cDNA was subsequently obtained by screening a mouse testis λ cDNA library followed by 5'RACE, as described under *Materials and Methods*. Its nucleotide and deduced amino acid sequences confirm that we cloned a novel NEP-like protein, referred to

thereafter as NL1.

NL1 cDNA spans 2925 nt, including a 5'-untranslated region of 331 nt, an open reading frame of 2295 nt from nt 332 to nt 2626, and a 3'-untranslated region of 299 nt. The sequence surrounding the proposed initiator ATG conforms to the Kozak consensus (Kozak 1986). The deduced amino acid sequence of NL1 reveals a putative type II transmembrane protein of 765 amino acid residues encompassing a short N-terminal cytoplasmic tail, a unique putative transmembrane domain, and a large C-terminal extracellular domain. The ectodomain contains nine potential N-glycosylation sites (Asn-X-Ser/Thr) and ten cysteine residues corresponding to those conserved among all the members of the family, which are presumably involved in proper folding and in maintenance of the protein in an active conformation. All amino acid residues known to be part of the active site of NEP are present in NL1. The predicted protein presents greater similarities to NEP than to any other member of the family.

Although NL1 shares numerous features with proteins of the neprilysin family, a notable aspect distinguishes it from the others : the first conserved cysteine residue of the ectodomain is more distant (34 amino acid residues) from the predicted transmembrane domain in NL1 than it is in NEP (9 residues) or any other members of the family. Moreover, we noticed a putative furin cleavage site (-Arg₅₈-Thr-Val-Val-Lys-Arg₆₃-) between the end of the transmembrane domain and the first cysteine. This observation suggests that NL1 could exist as a secreted rather than a membrane-bound protein.

NL1 expression in HEK 293 cells

HEK 293 cells were transfected with pCDNA3/RSV expression vector containing NL1 cDNA, and a permanent cell line was established as described under *Materials and Methods* (HEK/NL1 cells). Immunoblotting with a polyclonal antibody showed that after 16h of culture, most NL1 was present in the culture medium with small amounts of the enzyme in the cell extract. Secreted and cell-associated NL1 had apparent molecular masses of approximately 125 and 110 kDa, respectively. To characterize the glycosylation state of NL1, we next submitted the recombinant protein to deglycosylation by peptide : N-glycosidase F (PNGase F) and endoglycosidase H (endo H). PNGase F removes high mannose as well as most complex N-linked oligosaccharides added in the Golgi complex. In contrast, endo H removes N-linked oligosaccharide side chains of the high mannose type found on proteins in the RER but which have not yet transited through the Golgi complex; thus, resistance to endo H can be used as an indication that the protein has traveled through the Golgi complex. PNGase F treatment showed that the cell-associated and secreted NL1 were N-glycosylated as their electrophoretic mobility increased following digestion. However, the secreted NL1 migrated as a doublet after PNGase F treatment, with one

band co-migrating with cell-associated form and the second having a slower rate of migration. Since untreated and endo H-digested secreted NL1 are seen as single bands by SDS-PAGE, our observation suggests that a proportion of secreted NL1 undergoes further post-RER postranslational modification that renders some of the N-linked oligosaccharides resistant to PNGase F digestion.

In contrast to secreted NL1, NL1 from cell extract was sensitive to endo H treatment. This result shows differences in the glycosylation state of the two species and suggests that the cell-associated form observed in transfected cells is an intracellular species that has not traveled through the Golgi complex.

Processing of NL1 by a subtilisin-like convertase

To determine whether a member of the mammalian subtilisine-like convertase family is responsible for NL1 presence in the culture medium of transfected cells, we co-transfected transiently HEK 293 cells with a constant amount of plasmid pCDNA3/RSV/NL1 and increasing amounts of plasmid pCDNA3/CMV/PDX (Benjannet 1997). This latter vector promotes the expression of the α 1-antitrypsin Portland variant, α 1-PDX, a known inhibitor of subtilisin-like convertases (Anderson 1993). Immunoblot analysis of the culture media of cells expressing both NL1 and α 1-PDX indicated that NL1 secretion was strongly inhibited by the presence of α 1-PDX: a relation was observed between the amounts of α 1-PDX and the level of inhibition of NL1 secretion.

To confirm that proteolysis by the subtilisin-like convertase occurred at the putative furin cleavage site identified in NL1 ectodomain (-Arg₅₈-Thr-Val-Val-Lys-Arg₆₃-), the amino acid residues Asn₆₂-Gly₆₃ were substituted for Lys₆₂-Arg₆₃ by site-directed mutagenesis in vector pCDNA3/RSV/NL1 and the mutated vector used to establish HEK 293 cells expressing the mutant protein (HEK/NL1mut cells). Immunoblot analysis of the culture media of HEK/NL1mut cells showed that the mutation totally abolished secretion of NL1. Furthermore, an additional form of NL1 with a molecular mass of 127 kDa was detected in the extract of these cells. This new species was resistant to endo H digestion and was found associated with membranes when HEK/NL1mut cells were fractionated according to Chidiac *et al.* 1996 (result not shown).

NL1 enzymatic activity

Culture media from HEK 293 and HEK/NL1 cells were tested for enzymatic activity using as substrate (Tyrosyl-[3,5-³H])(D-Ala₂)-Leu₅-enkephalin, a known NEP substrate. Activity was detected in the culture medium of HEK/NL1 cells but not in that of HEK 293 cells. This activity increased linearly with the amounts of NL1 and with the incubation period, indicating that degradation of the substrate was due to NL1 enzymatic activity.

We next characterized NL1 enzymatic parameters using the same substrate

and compared them to those of an engineered soluble form of NEP (sNEP) (Lemay 1989). NL1 affinity for D-Ala₂-Leu₅-enkephalin was slightly higher than that of sNEP as shown by their K_m values of 18 μ M and 73 μ M, respectively. Inhibition assays showed that phosphoramidon had similar effects on NL1 and sNEP activity, with IC_{50} values of 0.9 nM and 0.5 nM respectively, and that thiorphan, a specific inhibitor of NEP, inhibited NL1 with an IC_{50} of 47 nM, as compared with an IC_{50} of 8 nM for NEP.

Very low levels of phosphoramidon-sensitive activity was detected in extracts of HEK/NL1 cells (data not shown) consistent with the small amounts of NL1 observed by immunoblotting.

To determine whether NL1 had cleavage site specificity similar to NEP, we incubated Leu₅-enkephalin in the presence of NL1 recovered from the medium of HEK/NL1 cells or in the presence of sNEP, and analyzed the degradation products by RP-HPLC. Peaks co-migrating with standard Tyr-Gly-Gly and Phe-Leu peptides were observed in both RP-HPLC profiles, indicating that both enzymes cleaved the substrate at the Gly₃-Phe₄ peptide bond. This enkephalin-degrading activity was completely inhibited by phosphoramidon (1 μ M).

Tissue and cellular distribution of NL1 mRNA

Tissue distribution of NL1 mRNA was determined by Northern blot analysis with a specific probe corresponding to the 5' end of the coding region of NL1 cDNA. A single transcript of 3.4 kb was detected exclusively in testis among all the mouse tissues tested. Mouse tissues were also screened by RT-PCR. Using this more sensitive technique, expression of NL1 was observed in several other tissues including heart, brain, spleen, lungs, liver and kidney. Consistent with the Northern blot results, RT-PCR analysis, although not strictly quantitative, detected more NL1 mRNA in testis than in other tissues.

To gain more insight into NL1 mRNA distribution, we examined by ISH cryostat sagittal sections from a 4-day newborn mouse, as well as sections from a 16-day old animal (p16) and adult tissues (heart, brain, spleen, lungs, liver, kidney and testis). The presence of NL1 mRNA was detected only in adult testis. Only the germinal cells in the luminal face of the seminiferous tubules were labeled. These cells were identified as round and elongated spermatids in all spermiogenesis maturational stages. Neither spermatozoa nor spermatocytes, spermatogonies or Sertoli cells were labeled. Interstitial cells were also negative. Controls were performed with sense riboprobes, which produced only nonspecific background (data not shown). The 4-day old mouse sagittal sections and all other tissues tested were negative.

DISCUSSION

The great interest in members of the Neprilysin family as putative therapeutic

5 targets, and the recent discovery of new members of this important family of
peptidases led us to investigate whether additional members of the family remained
to be identified. Using a PCR-based strategy, we cloned, from mouse testis, a partial
cDNA encoding a new NEP-like enzyme that we called NL1. Analysis of the amino acid
10 sequence encoded by the full-length NL1 cDNA revealed that this member of the
family resembles NEP the most: 55% identity and 74% similarity. Recently, the primary
structure of a new zinc metallopeptidase from total mouse embryo was reported (Ikeda
1999). This enzyme, called SEP, is found either as a soluble or a cell-associated form
due to alternative splicing. NL1 shows only 3 amino acid differences with the soluble
15 form of SEP indicating that secreted SEP and NL1 are the same enzyme. Our cloning
strategy did not allow characterization of the cell-associated form of NL1 which is a
minor species in mouse testis (Ikeda 1999).

The amino acid sequence of NL1 predicts a topology of a type II integral membrane
glycoprotein that is similar to the other members of the family. Treatment of the
20 recombinant protein with PNGase F showed that indeed NL1 possesses N-linked
carbohydrate side chains. However, it is not possible to determine precisely whether
all nine putative N-glycosylation sites are used, but the 30 kDa decrease in molecular
mass upon PNGase F treatment suggests that most are glycosylated. It has already
been shown that all asparagine residues in a Asn-X-Ser/Thr consensus are
25 glycosylated in rabbit NEP expressed in COS-1 cells and that sugar moieties increase
the stability and enzymatic activity of the protein and facilitate its intracellular transport
(Lafrance 1994). Three of NEP glycosylated Asn residues (Asn 145, Asn 285 and Asn
628) are conserved in NL1 (Asn 163, Asn 303 and Asn 643). Amongst these residues,
Asn 145 and Asn 628 have been reported to influence NEP enzymatic activity
30 (Lafrance 1994). In the same work, it has also been shown that the effect of sugar
addition on folding and intracellular transport of NEP is a cumulative effect of all
glycosylation sites rather than a contribution of any particular one. Glycosylation of
NL1 may share similarities with that of NEP since we found their primary structures
and enzymatic activities to be very similar.

35 Surprisingly, expression of the cDNA by transfection of HEK 293 cells showed that
most of the enzyme was secreted in the culture medium. The small amount of NL1
associated with the cells was endo H-sensitive, suggesting that the cell-associated
enzyme is a species that has not yet left the RER. The presence of a furin cleavage
site in NL1 sequence between the predicted transmembrane domain and the first
conserved cysteine residue of the ectodomain led us to believe that a member of the
mammalian subtilisin-like family of convertases was responsible for the presence of
NL1 in the culture medium. These enzymes are involved in processing a variety of
precursor proteins such as growth factors and hormones, receptors, plasma proteins,
matrix metalloproteinases, metalloproteases-desintegrins and viral envelope

glycoproteins [for a review see: (Nakayama 1997). Site-directed mutagenesis of the furin cleavage site (-Arg₅₈-Thr-Val-Val-Lys-Arg₆₃-) and expression of α 1-PDX, a potent inhibitor of mammalian subtilisin-like convertases (Anderson 1993), confirmed that a member of this family of endoproteases was involved in NL1 secretion presumably by

5 cleaving in carboxy-terminus of Arg₆₃. There are only a few examples of proteins which are processed from a membrane-bound precursor to a secreted form following cleavage by subtilisin-like convertases; these include meprin and collagen XVII (Milhiet 1995; Schacke 1998). Three members of the subtilisin-like family of convertases, namely furin, PC4 and PC7, are known to be expressed in germ cells (Nakayama

10 1992; Torri 1993; Seidah 1992, 1996). Whether one of these convertases generates secreted NL1 from its membrane form is under current investigation. In any case, NL1 is the only known member of the neprilysin family that is secreted. This unique feature suggests that NL1 plays its physiological role in a context different from that of the membrane-bound peptidases, thereby diversifying the role of the peptidases of the

15 neprilysin family. It is of interest that circulating forms of NEP in blood and urine have been described, but they have generally been related to pathological or stressful conditions (Almenoff 1984; Deschodt-Lanckmann 1989; Johnson 1985; Soleilhac 1996; Aviv 1995).

We have observed in cells expressing NL1 mutated at the furin cleavage site the

20 appearance of a species resistant to digestion by endo H. This mutated protein was associated with cellular membranes. Taken together, these results indicate that NL1 is first synthesized and inserted in the RER membrane as a type II transmembrane protein. During intracellular transport, NL1 is converted to a soluble form by the action of a member of the mammalian subtilisin-like convertases. The identity of the cellular

25 compartment where this process occurs is not known. However, mammalian subtilisin-like convertases are usually active in post-Golgi compartments of the secretory pathway suggesting that processing of NL1 from the membrane bound form to the soluble form is a post-Golgi event.

Despite almost total abrogation of NL1 secretion, we observed only a slight

30 accumulation of endo H-resistant NL1 in cells either co-expressing α 1-PDX and NL1 (result not shown) or expressing mutated NL1. This observation suggests that unprocessed NL1 is rapidly degraded. A similar behavior was reported for the Notch1 receptor expressed in the furin-deficient cell line LoVo (Logeat 1998). The mechanism(s) by which these unprocessed proteins are degraded is still unknown. It

35 is interesting to point out that the spliceform of SEP that has lost a 23 amino acid peptide, including the furin cleavage site, generates a cell-associated endo H-sensitive molecule (Ikeda 1999).

The most important observation regarding the NL1 primary structure is the conservation of residues which in NEP are essential for catalysis and binding of

substrates or inhibitors. This finding suggests that NL1 could effectively act as an endopeptidase with a catalytic mechanism similar to that of NEP. This hypothesis was supported by the demonstration that D-Ala₂-Leu₅-enkephalin, a peptide substrate often used to monitor NEP activity, was also an excellent NL1 substrate. The affinity of NL1 for D-Ala₂-Leu₅-enkephalin was even higher than that of NEP, as reflected by a K_m value 4- to 5-fold lower. Furthermore, two well known NEP inhibitors, phosphoramidon and thiorphan, also abolished NL1 activity. Phosphoramidon, which inhibits NEP as well as ECE-1 activity, albeit to a lesser extent (Turner 1996), had very similar effects on NL1 and NEP, with an IC₅₀ value for NL1 varying not more than two-fold from the value determined for NEP. Thiorphan, considered to be a more specific inhibitor of NEP, also inhibited NL1 activity, with an IC₅₀ six-fold greater than that for NEP. These results suggest that the active sites of NL1 and NEP are similar. This hypothesis is supported by the observation that secreted SEP degraded a set of peptides known to be NEP substrates, including substance P, bradykinin and atrial natriuretic peptide (Ikeda 1999). Taken together, these results illustrate the importance of identifying and characterizing other member of the family for the design of highly specific inhibitors.

In agreement with the enzymatic parameters demonstrating that NL1 and NEP have similar catalytic sites, we have observed that both enzymes cleaved Leu₅-enkephalin at the same peptide bond. This result suggests that NL1 hydrolyzes peptide bonds on the amino side of hydrophobic amino acid residues as does NEP (Turner 1985). However, several other peptides will have to be tested to confirm this specificity and to determine whether NL1 has dipeptidyl carboxypeptidase activity as was shown for NEP (Malfroy 1982; Bateman 1989; Beaumont 1991) and more recently for ECE-1 (Johnson 1999).

RT-PCR experiments with specific primers for the soluble and cell-associated forms of SEP showed a wide tissue distribution of the enzyme with the soluble form of SEP being predominant in testis and the cell-associated form in other tissues (Ikeda 1999). Our RT-PCR results confirmed the wide tissue distribution of NL1. However, Northern blotting and *in situ* hybridization experiments indicated that expression of NL1 is predominant in germ cells of mature testis. Interestingly, proenkephalin mRNA has been shown to be expressed in germ cells and somatic cells of the testis (Torii 1993, Seidah 1992; Kew 1989; Mehta 1994; Kilpatrick 1986, 1987). Specific functions for testicular enkephalin peptides have not yet been defined, but it is believed that they could act as intratesticular paracrine/autocrine factors. In addition to their putative role as mediators of testicular cell communication, it has also been demonstrated that proenkephalin products synthesized by spermatogenic cells during spermatogenesis are stored in the acrosome of human, hamster, rat and sheep spermatozoa and are release from sperm following acrosomal reaction (Kew 1990). It has thus been proposed that proenkephalin products may act as sperm acrosomal factors during the

TOP SECRET

fertilization process as well as intratesticular regulators secreted by spermatogenic cells. Since Leu₅-enkephalin was found to be a good substrate for NL1, opioid peptides originating from proenkephalin could serve as physiological substrate for this new enzyme. In this way, NL1 would serve to regulate the activity of these bioactive peptides.

Testis is the only tissue where the soluble form of SEP is predominant (Ikeda, 1999), suggesting a testis-specific alternative splicing. Expression of testis-specific molecular species of peptidases or prohormones, arising through diverse mechanisms, has been documented in the past (Howard 1990; Jeannotte 1987). However, the physiological significance of these testis-specific species is not always clear. In the case of NL1 or SEP, it might allow local constitutive secretion by germinal cells of an otherwise cell-associated enzyme, to regulate spermatogenesis much like several other proteolytic enzymes of the seminiferous tubules (Monsees 1998). Alternatively, it might allow accumulation in acrosome with proenkephalin peptides and release upon acrosomal reaction. More exhaustive studies concerning NL1 localization and physiological substrate identification will be needed to understand its role in the testis and possibly in the fertilization process.

Cloning of other members of the family

To find other members of the NEP-like family, we will use the same RT-PCR strategy to amplify mRNA isolated from tissues known to be regulated by peptidergic systems (brain, thymus, kidney, heart, lung, ovary, pancreas, bone, bone marrow and lymphoid cells). In fact, many of these tissues are known to express at least one member of the family and/or to control a peptidergic pathway on which peptidase inhibitors have major effects. Amplified fragments will be cloned and the resulting clones will be sequenced and compared to the sequence of known peptidases, as described above. Pairs of degenerate primers in other highly conserved regions will also be designed to increase the possibility of cloning other relevant peptidases.

DISCUSSION

As discussed above, peptidases of the NEP family known to date have often been found to play important physiological roles. This is certainly true for NEP itself, ECEs and PEX, (see review above). For this reason, some of these enzymes (as it was the case for NEP and ECE in the past) might be interesting targets for the design of inhibitors that in turn could be used as therapeutic agents in various pathological conditions. However, it is of some concern that inhibitors designed for one enzyme may also inhibit to some extent other members of the family. This lack of specificity for an inhibitor used as a therapeutic agent in the long term treatments such as those used as antihypertensive agents for instance, may cause unforeseen problems due to unwanted side effects. The objectives of the present work was to develop a strategy

to clone new members of the NEP family of peptidases. The results presented in this report clearly show that our strategy can be successful. We have determined the complete or partial nucleotide sequence of three cDNAs encoding putative enzymes of the NEP family.

5 These cDNA sequences are valuable tools and may be used to:

Produce antibodies

10 As shown in the present work, knowledge of NL cDNA sequences can be used to raise specific antibodies. For example but not exclusively, regions of less homology between the peptidases (amino acid residues 50 to 450) can be used to synthesize peptides whose sequences are deduced from the translation of the cDNAs, and/or bacterially-expressed fragments of the cDNAs fused for example but not exclusively to GST may be purified and injected into rabbits or mice for polyclonal or monoclonal antibody production. These antibodies can be used to:

- 15 - identify by immunohistochemistry the peptidergic pathways in which the peptidases are functioning;
- 20 - study the physiopathology of NL-enzymes by immunoblotting or immunohistochemistry on samples of biological fluids or biopsies;
- 25 - set up high through put screening assays to identify NL-enzymes inhibitors. This can be done for example but not exclusively by using the antibodies to attach the NL-enzymes to a solid support;
- 30 - purify NL-enzymes with said antibodies by immunoprecipitation or affinity chromatography by identifying antibodies capable of selectively binding to the NL-enzymes in one set of conditions and releasing it in another set of conditions typically involving a large pH or salt concentration change without denaturing the NL-enzyme;
- identify antibodies that block NL-enzymes activities and use them as therapeutic agents. Blocking antibodies can be identified by adding antisera or ascite fluid to an *in vitro* enzymatic assay and looking for inhibition of NL-enzymes activities. Blocking antibodies could then be injected to normal or disease model animals to test for *in vivo* effects.

Derive specific RNA or DNA probes

35 As shown in the present work, knowledge of the nucleotide sequence of the members of the NEP-family allows nucleotide sequence comparisons and facilitate the design of specific RNA or DNA probes by methods such as but not exclusively molecular cloning, *in vitro* transcription, PCR or DNA synthesis. The probes thus obtained can be used to:

- derive specific probes or oligonucleotides for RNA and DNA analysis, such as Northern blot and *in situ* hybridization, chromosome mapping

FOR "SECRET" 660

or PCR testing. These probes could be used for genetic testing of normal or pathological samples of biological fluids or biopsies;

- make vectors for gene knock-out or knock-in in mice. The long range PCR technique and/or screening of a mouse genomic library with probes derived from the 5'-end of the cDNAs can be used to isolate large exon/intron regions. We will then substitute one or more of the cloned genomic DNA exons for the neomycin resistance expression cassette for producing homologous recombination and knock-out mice.

Alternatively, cDNAs coding for NLs will be used to overexpressed each of these enzymes in transgenic mice. The cDNAs will be cloned downstream from a promoter sequence, and injected in fertilised mouse eggs. Depending on specific questions to be answered, the chosen promoter sequence will allow expression of the peptidases either in every tissues or in a cell- or tissue-specific manner. Injected eggs will be transferred into foster mothers and the resulting mice analysed for peptidase expression;

- replace defective NL genes in a gene therapy strategy. The NL full length cDNAs could be cloned under the control of a constitutive or inducible promoter having a narrow or wide range of tissue expression and introduced with appropriate vectors in subjects having defective genes;

- synthesise oligonucleotides that could be used to interfere with the expression of the NLs. For example but not exclusively, oligonucleotides with antisens or ribozyme activity could be developed. These oligonucleotides could be introduced in subjects as described above;

- isolate other members of the family. Screening cDNA and/or genomic libraries with these cDNA probes at low stringency may allow to clone new members of the NEP-like family. Alternatively, alignment of the sequences may allow one to design specific degenerate oligonucleotide primers for RT-PCR screening with mRNA from tissues such as but not exclusively, the hearth and the brain.

Production of recombinant NL-enzymes

As shown in the present work, recombinant active NL-enzymes can be obtained by expression of NL-cDNAs in mammalian cells. From past experience with neprilysin, another member of the family (Devault *et al.*, 1988; Fossiez *et al.*, 1992; Ellefsen *et al.*, submitted), expression can also be performed in other expression systems after cloning of NL-cDNAs in appropriate expression vectors. These expression systems may include but not exclusively the baculovirus/insect cells or larvae system and the *Pichia*

FOR "SECRETED"

5

10

15

20

25

30

35

pastoris-based yeast system. Production of recombinant NL-enzymes includes the production of naturally occurring membrane bound or soluble forms of the proteins or genetically engineered soluble forms of the enzymes. The latter can be obtained by substituting the cytosolic and trans-membrane domain by a cleavable signal peptide such as that of proopiomelanocortin, but not exclusively, as done previously (Lemay *et al.*, 1989) or by transforming by genetic manipulations the non-cleavable signal peptide membrane anchor domain into a cleavable signal peptide, as done previously (Lemire *et al.*, 1997) or by fusion of the ectodomain of NL-enzymes to the amino-terminal domain (from the initiator methionine to amino acid residue 300) of naturally occurring soluble NLs such as, but not exclusively, NL-1 as done in this work.

These recombinant NLs could be used to:

- find a substrate. A substrate can be identified using one of the following.
- Screening of existing bioactive peptides. Peptides are incubated in the presence of NL-enzymes and subsequently analysed by HPLC for degradation. Degradation is observed by disappearance of the peak of substrate and the appearance of peaks of products;
- Screening phage libraries specifically designed for the purpose (phage display library). Each phage expresses at its surface, as part of its coat protein, a random peptide sequence preceded by a peptide sequence recognisable by an antibody or any other sequence-recognizing agent. This latter sequence serves to attach the phage to a solid support. Upon addition of the NL-enzyme the random sequences that are NL substrate are cleaved, releasing the phage. After several rounds of cleavage, the phage sequence is determined to identify the peptide segment recognized by the enzyme.
- Extract of the tissue where the enzyme is expressed is collected and prepared for chromatographic analysis (HPLC, capillary electrophoresis or any other high resolution separation system) by denaturing the extracted proteins with a solvent (acetonitrile or methanol). The extract is subjected to chromatographic separation. The same extract is incubated with the enzyme for a period sufficient to observe a difference between the 2 chromatograms. The regions with the identified changes are collected and subjected to mass spectrometric analysis to determine the peptide compositions.
- Small peptide libraries are prepared with a fluorophore at one extremity and a quencher group at the other (Meldal *et al* *Methods in molecular biology* 1998,87). The substrate can be identified using a strategy described in Apletalina *et al* (*JBC* (1998)273, 41, 26589-95). For each

hexapeptide library, the identity of one residue at one position remains constant while the rest is randomized (for a total of $6 \times 20 = 120$ individual libraries). Each library is made-up of 3.2 million different members and is identified both by the position of the constant residue along the hexapeptide, and its identity. The NL-enzyme is added to each library and the fluorescence is recorded. The data is organized to identify the libraries producing the most fluorescence for each position along the hexapeptide. This arrangement suggests the identity of important residues at each position along the hexapeptide. Hexapeptide representing the best suggestions are prepared and tested in a similar fashion. From this set, the hexapeptide with the best fluorescence is selected.

set up enzymatic assays. An enzymatic assay consists in the addition of the above-identified substrate to the enzyme in constant conditions of pH, salts, temperature and time. The resulting solution is assayed for the hydrolysed peptide or for the intact peptide. This assay can be realized with specific antibodies, HPLC or, when self-quenched fluorescence tagged peptides are used (Meldal et al), by the appearance of fluorescence. The enzyme may be in solution or attached to a solid substrate;

identify inhibitors. Inhibitors can be identified from synthetic libraries, biota extracts and from rationally designed inhibitors using X-ray crystallography and substituent activity relationships. Each molecule or extract fraction is tested for inhibitory activity using the enzymatic test described above. The molecule responsible for the largest inhibition is further tested to determine its pharmacological and toxicological properties following known procedures. The inhibitor with the best distribution, pharmacological action combined with low toxicity will be selected for drug manufacturing. Pharmaceutically acceptable formulation of the inhibitor or its acceptable salt will be prepared by mixing with known excipients to produce tablets, capsules or injectable solutions. Between 1 and 500mg of the drug is administered to the patients;

inject the native or soluble purified NL-enzymes into subjects. In the case of disease or pathologies caused by a lack or decrease in NL activity, the purified NL could be injected intravenously or otherwise in patients. Alternatively, immobilized NL-enzymes could be introduced at the site of orthopedic surgery or implantation of devices in bones or dental tissues.

FOR OFFICIAL USE ONLY

Secretion of foreign proteins and peptides

As shown in the present work, the amino-terminal domain of NL-1 (from the initiator methionine to the furin site) can be used to promote the secretion of a foreign protein (in this case the ectodomain of NL-3 and β -endorphin).

The amino-terminal domain of NL-1 but also of other naturally occurring soluble NL-enzymes could be used to:

- promote production and secretion of foreign proteins. This can be achieved by genetically fusing sequences coding for said foreign proteins downstream from and in phase with the amino-terminal of NL-1. These chimeric constructs could be introduced with the help of appropriate vectors in any of the expression systems mentioned above for protein production and secretion;
- promote production and secretion of bioactive peptides. Sequences encoding small bioactive peptides such as but not exclusively β -endorphin, the enkephalins, substance P, atrial natriuretic peptide (ANF) and osteostatine, could be fused immediately downstream and in phase the furin site of NL-1. These DNA constructs could be used as described above to produce bioactive peptides.
- serve as model to design artificial (non-naturally occurring) proteins or protein segments (protein vectors) to promote secretion of proteins or peptides. These protein vectors can be constructed to resemble a secreted protein. In this case they would be assembled of an endoplasmic reticulum signal peptide, a spacer of varying length and a furin cleavage site to which the protein or peptide destined for secretion can be fused. The total length of the spacer, furin cleavage site and protein or peptide destined for secretion must be at least 70 amino acid residues. Alternatively, such protein vectors could be assembled to resemble a type II membrane protein. In this case they would comprise from the amino to the carboxy-terminus a cytosolic domain of varying length, a transmembrane domain that also acts as a signal peptide, an extracellular segment of varying length and a furin cleavage site to which the protein or peptide destined for secretion can be fused. The total length of the extracellular segment, furin cleavage site and protein or peptide destined for secretion must be at least 70 amino acid residues.

Therapeutic applications of NL-enzymes

The inappropriate processing of endogenous peptides causes several diseases. The inappropriate processing may result from pathologic

FOOTNOTES

concentration of the enzyme itself, its substrate or other elements of the biochemical machinery downstream from the controlling enzyme. In this context it is possible to help the patient by managing the activity of the controlling enzyme.

- 5 - NL-enzymes have been localized to the brain and may be involved in the improper processing of β -amyloid precursor. Inhibitions of this process by drugs prepared as above, will help patients with Alzheimer disease as well as other patient suffering from diseases caused by plaque formation;
- 10 - NL-enzymes may be involved in the improper processing of other peptides involved in neurological diseases, pain or psychiatric disorders. Appropriately designed inhibitors will help in the management of such diseases;
- 15 - NL-1 is found in testis and is associated with spermatozoid maturation. Peptides improperly processed by the enzyme may lead to infertility. The addition of NL-1 ex-vivo to seminal liquid or immature spermatozoids taken directly from testis during an in-vitro fertilization procedure will increase fertility. Conversely, the use of a small-molecule inhibitor or removal of NL-1 with an antibody could increase fertility during an in-vitro fertilization procedure. The administration of a NL-1 inhibitor may increase or decrease the fertility potential. This inhibitor is formulated and administered as described above.
- 20 - NL-3 is found in ovaries and may be involved in the processing of a peptide involved in the maturation of eggs. The addition of NL-3 ex-vivo to immature eggs taken directly from ovaries during an in-vitro fertilization procedure will increase fertility. Conversely, the use of a small-molecule inhibitor or removal of NL-3 with an antibody could increase fertility during an in-vitro fertilization procedure. This inhibitor is formulated and administered as described above;
- 25 - NL-3 is found in bones. The improper processing of peptides by the enzyme may result in bone disease or abnormal phosphate metabolism. Administration of an inhibitor, as described above, will allow the disease management.
- 30
- 35

TABLE I

Extend of amino acid sequence identity between members of the NEP-like family

	hNEP	hPEX	hECE-1A	hECE-2	hKELL	sNL-1	hNL-2	hNL-3
hNEP	100*							

5

hPEX	35	100						
hECE-1A	39	38	100					
hECE-2	36	37	62	100				
hKELL	23	24	30	31	100			
sNL-1	55	39	39	39	26	100		
hNL-2	54	39	39	39	26	77	100	
hNL-3	35	32	37	37	28	36	34	100

*: percentage of sequence identity

FOIA b 7 - D

REFERENCE LIST

- Almenoff, J., Teirstein, A.S., Thornton, J.C., and Orlowski, M. (1984) *J. Lab. Clin. Med.* **103**, 420-431
- Anderson, A.D., Thomas, L., Hayflick, J.S., and Thomas, G. (1993) *J. Biol. Chem.* **268**,
5 24887-24891
- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., and Smith, J.A. (1988). *Current protocols in molecular biology* (New York: Wiley Interscience).
- Aviv, R., Gurbanov, K., Hoffman, A., Blumberg, S., and Winaver, J. (1995) *Kidney Int.* **47**, 855-860
- 10 Bateman, R.C., Jackson D., Slaughter, C.A., Unnithan, S., Chai, Y.G., Moomaw, C.,
and Hersh, L.B. (1989) *J. Biol. Chem.* **264**, 6151-6157
- Bawab, W., Aloyz, R.S., Crine, P., Roques, B.P., DesGroseillers, L. (1993) *Biochem. J.* **296**, 459-465
- Beaumont, A., Le Moual, H., Boileau, G., Crine, P., and Roques, B.P. (1991).
15 Evidence that both arginine 102 and arginine 747 are involved in substrate
binding to neutral endopeptidase (EC 3.4.24.11). *J.Biol.Chem.* **266**, 214-220.
- Benjannet, S., Savaria, D., Laslop, A., Munzer, J.S., Chrétien, M., Marcinkiewicz, M.,
and Seidah, N.G. (1997) *J. Biol. Chem.* **272**, 26210-26218.
- Chen, C. and Okayama, H. (1987) *Mol. Cell Biol.* **7**, 2745-2752
- 20 Chidiac, P., Nouet, S., and Bouvier, M. (1996) *Mol. Pharmacol.* **50**, 662-669
- Corvol, P. and Williams, T.A. (1997). *Biochemical and molecular aspects of
angiotensin I-converting enzyme. In Cell-surface peptidases in health and
disease. A.J. Kenny and C.M. Boustead, eds. (Oxford, UK: BIOS Scientific
Publishers Ltd), pp. 99-117.*
- 25 Crine, P., Dion, N., and Boileau, G. (1997). *Endopeptidase-24.11. In Cell-Surface
Peptidases in Health and Disease. A.J. Kenny and C.M. Boustead, eds.
(Oxford: BIOS Scientific Publishers), pp. 79-98.*
- Crine, P., LeGrimellec, C., Lemieux, E., Labonté, L., Fortin, S., Blachier, A. and Aubry,
M. (1985) The production and characterization of a monoclonal antibody
30 specific for the 94000 dalton enkephalin-degrading peptidase from rabbit kidney
brush border. *Biochem. Biophys Res. Commun.* **131**, 255-261.
- Deschodt-Lanckman, M., Michaux, F., de Prez, E., Abramowicz, D., Vanherweghem,
J.L., and Goldman, M. (1989) *Life Sci.* **45**, 133-141
- Devault, A., Nault, C., Zollinger, M., Fournié-Zaluski, M.-C., Roques, B.P., Crine, P.,
35 and Boileau, G. (1988a). Expression of neutral endopeptidase (enkephalinase)
in heterologous COS-1 cells. Characterization of the recombinant enzyme and
evidence for a glutamic acid residue at the active site. *J.Biol.Chem.* **263**, 4033-
4040.
- Devault, A., Sales, V., Nault, C., Beaumont, A., Roques, B.P., Crine, P., and Boileau,

- G. (1988b). Exploration of the catalytic site of endopeptidase 24.11 by site-directed mutagenesis. Histidine residues 583 and 587 are essential for catalysis. *FEBS Lett.* **231**, 54-58.
- 5 Dion, N., Le Moual, H., Crine, P., and Boileau, G. (1993). Kinetic evidence that His-711 of neutral endopeptidase 24.11 is involved in stabilization of the transition state. *FEBS Lett.* **318**, 301-304.
- Dion, N., Le Moual, H., Fournié-Zaluski, M.C., Roques, B.P., Crine, P., and Boileau, G. (1995). Evidence that Asn⁵⁴² of neprilysin (EC 3.4.24.11) is involved in binding of the P₂' residue of substrates and inhibitors. *Biochem.J.* **311**, 623-627.
- 10 Holmes, M.A. and Matthews, B.W. (1982). Structure of thermolysin refined at 1.6 Å resolution. *J.Mol.Biol.* **160**, 623-639.
- Hooper, N.M. (1994). Families of zinc metalloproteases. *FEBS Lett.* **354**, 1-6.
- Howard, T.E., Shai, S.Y., Langford, K.G., Martin, B.M., Bernstein, K.E. (1990) *Mol. Cell. Biol.* **10**, 4294-4302
- 15 Ikeda, K., Emoto, N., Raharjo, S.B., Nurhantari, Y., Saiki, K., Yokoyama, M., Matsuo, M. (1999) *J. Biol. Chem.* **274**, 32469-32477
- Jeannotte, L., Burbach, J.P., Drouin, J. (1987) *Mol. Endocrinol.* **1**, 749-757
- Johnson, A.R., Coalson, J.J., Ashton, J., Larumbide, M., and Erdos, E.G. (1985) *Am. Rev. Respir. Dis.* **132**, 1262-1267
- 20 Johnson, G.D., Stevenson, T., and Ahn, K. (1999) *J. Biol. Chem.* **274**, 4053-4058
- Endocrinol.* **3**, 1191-1196
- Kenny, J. (1993). Endopeptidase-24.11: Putative substrates and possible roles. *Biochem.Soc.Trans.* **21**, 663-668.
- Kew, D., Jin, D.F., Kim, F., Laddis, T., and Kilpatrick, D.L. (1989) *Mol. J. Biol Chem.* **266**, 214-220
- 25 Kew, D., Muffly, K.E., and Kilpatrick, D.L. (1990) *Proc. Natl. Acad. Sci. U.S.A* **87**, 9143-9147
- Kilpatrick, D.L., Borland, K., and Jin, D.F. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 5695-5699
- 30 Kilpatrick, D.L. and Millette, C.F. (1986) *Proc. Natl. Acad. Sci. U.S.A* **83**, 5015-5018
- Kozak, M. (1986) *Cell* **44**, 283-292
- Lafrance, M.H., Vezina, C., Wang, Q., Boileau, G., Crine, P., and Lemay, G. (1994) *Biochem. J.* **302**, 451-454.
- Lanctôt, C., Fournier, H., Howell, S., Boileau, G., and Crine, P. (1995) *Biochem. J.* **305**, 165-171
- 35 Le Moual, H., Devault, A., Roques, B.P., Crine, P., and Boileau, G. (1991). Identification of glutamic acid 646 as a zinc-coordinating residue in endopeptidase-24.11. *J.Biol.Chem.* **266**, 15670-15674.
- Le Moual, H., Dion, N., Roques, B.P., Crine, P., and Boileau, G. (1994). Asp650 is

- crucial for catalytic activity of neutral endopeptidase 24-11. *Eur.J.Biochem.* 221, 475-480.
- Lemay, G., Waksman, G., Roques, B.P., Crine, P., and Boileau, G. (1989). Fusion of a cleavable signal peptide to the ectodomain of neutral endopeptidase (EC 3.4.24.11) results in the secretion of an active enzyme in COS-1 cells. *J.Biol.Chem.* 264, 15620-15623.
- Logeat, F., Bessia, C., Brou, C., LeBail, O., Jarriault, S., Seidah, N.G., and Israël, A. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 8108-8112.
- Malfroy, B., and Schwartz, J.C. (1982) *Biochem. Biophys. Res.Com.* **106**, 276-285
- 10 Marie-Claire, C., Ruffet, E., Antonczak, S., Beaumont, A., O'Donohue, M., Roques, B.P., and Fournié-Zaluski, M.C. (1997). Evidence by site-directed mutagenesis that arginine 203 of thermolysin and arginine 717 of neprilysin (neutral endopeptidase) play equivalent critical roles in substrate hydrolysis and inhibitor binding. *Biochemistry* 36, 13938-13945.
- 15 Mehta, N.D., Don, J., Zinn, S.A., Millette, C., Wolgemuth, D.J., and Kilpatrick, D.L. (1994) *Endocrinology* **135**, 1543-1550
- Milhiet, P.E., Chevallier, S., Corbeil, D., Seidah, N.G., Crine, P., and Boileau, G. (1995) *Biochem. J.* **309**, 683-688
- Monsees, T.K., Gornig, M., Schill, W.B., and Miska, W. (1998) *Andrologia* **30**, 185-191
- 20 Nakayama, K. (1997) *Biochem. J.* **327**, 625-635
- Nakayama, K., Kim, W.S., Torii, S., Hosaka, M., Nakagawa, T., Ikemizu, J., Baba, T., and Murakami, K. (1992) *J. Biol. Chem.* **267**, 5897-5900
- Phelan, M.C., Rogers, R.C., Clarkson, K.B., Bowyer, F.P., Levine, M.A., Estabrooks, L.L., Severson, M.C., and Dobyns, W.B. (1995). Albright hereditary osteodystrophy and del(2) (q37.3) in four unrelated individuals. *Am.J.Med.Genet.* 58, 1-7.
- 25 Raut, R., Rouleau, J.L., Blais, C. Jr, Gosselin, H., Molinaro, G., Sirois, M.G., Lepage, Y., Crine, P., and Adam, A. (1999) Bradykinin metabolism in the postinfarcted rat heart: role of ACE and neutral endopeptidase 24.11. *Am. J Physiol.* 276, H1769-H1779.
- 30 Ruchon, A.F., Marcinkiewicz, M., Siegfried, G., Tenenhouse, H.S., DesGroseillers, L., Crine, P., and Boileau, G. (1998). Pex mRNA is localized in developing mouse osteoblasts and odontoblasts. *J.Histochem.Cytochem.* 46, 459-468.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular cloning, a laboratory manual* (New-York: Cold Spring Harbor Laboratory Press).
- 35 Schacke, H., Schumann, H., Hammami-Hauasli, N., Raghunath, M., and Bruckner-Tuderman, L. (1998) *J. Biol. Chem.* **273**, 25937-25943
- Seidah, N.G. and Chrétien, M. (1995). Pro-protein convertases of subtilisin/kexin family. *Methods Enzymol.* 244, 175-188.

- Seidah, N.G., Day, R., Hamelin, J., Gaspar, A., Collard, M.W., and Chretien, M. (1992) *Mol. Endocrinol.* **6**, 1559-1570
- Seidah, N.G., Hamelin, J., Mamarbachi, M., Dong, W., Tadros, H., Mbikai, M., Chrétien, M., and Day, R. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 3388-3393
- 5 Soleilhac, J.M., Lafuma, C., Porcher, J.M., Auburtin, G., and Roques, B.P. (1996) *Eur. J. Clin. Invest.* **26**, 1011-1017
- Torii, S., Yamagishi, T., Murakami, K., and Nakayama, K. (1993) *FEBS Lett.* **316**, 12-16
- 10 Turner, A.J. (1997a). Endothelin-converting enzymes. In *Cell-surface peptidases in health and disease*. A.J. Kenny and C.M. Boustead, eds. (Oxford, UK: BIOS Scientific Publishers Ltd.), pp. 137-153.
- Turner, A.J. and Tanzawa, K. (1997b). Mammalian membrane metallopeptidases: NEP, ECE, KELL, and PEX. *FASEB J.* **11**, 355-364.
- Turner, A.J., and Murphy, L. (1996) *Biochem. Pharmacol.* **51**, 91-102
- 15 Turner, A.J., Matsas, R., and Kenny, A.J. (1985) *Biochem. Pharmacol.* **34**, 1347-1356
- Vijayaraghavan, J., Kim, Y.-A., Jackson, D., Orłowski, M., and Hersh, L.B. (1990). Use of site-directed mutagenesis to identify valine-573 in the S'1 binding site of rat neutral endopeptidase 24.11 (enkephalinase). *Biochemistry* **29**, 8052-8056.

FOI 2006-0286